



Int J Environ Res Public Health. Apr 2009; 6(4): 1418–1442.

PMCID: PMC2681191

Published online Apr 14, 2009. doi: [10.3390/ijerph6041418](https://doi.org/10.3390/ijerph6041418)

## Perchlorate: Health Effects and Technologies for Its Removal from Water Resources

[Asha Srinivasan](#)<sup>1</sup> and [Thiruvengkatachari Viraraghavan](#)<sup>2,\*</sup>

<sup>1</sup> Faculty of Engineering, University of Regina, Regina, SK, S4S 0A2, Canada; E-Mail: [sriniash@uregina.ca](mailto:sriniash@uregina.ca)

<sup>2</sup> Professor Emeritus, Faculty of Engineering, University of Regina, Regina, SK, S4S 0A2, Canada

\* Author to whom correspondence should be addressed; E-Mail: [T.Viraraghavan@uregina.ca](mailto:T.Viraraghavan@uregina.ca); Tel.: +1-306-585-4094; Fax: +1-306-585-4855

Received February 13, 2009; Accepted April 3, 2009.

Copyright © 2009 by the authors; licensee Molecular Diversity Preservation International, Basel, Switzerland.

This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).

This article has been [cited by](#) other articles in PMC.

### Abstract

[Go to:](#)

Perchlorate has been found in drinking water and surface waters in the United States and Canada. It is primarily associated with release from defense and military operations. Natural sources include certain fertilizers and potash ores. Although it is a strong oxidant, perchlorate is very persistent in the environment. At high concentrations perchlorate can affect the thyroid gland by inhibiting the uptake of iodine. A maximum contaminant level has not been set, while a guidance value of 6 ppb has been suggested by Health Canada. Perchlorate is measured in environmental samples primarily by ion chromatography. It can be removed from water by anion exchange or membrane filtration. Biological and chemical processes are also effective in removing this species from water.

**Keywords:** Perchlorate, Drinking Water, Toxicity, Health Effects, Treatment Technologies

### 1. Introduction

[Go to:](#)

The perchlorate anion consists of a tetrahedral array of oxygen atoms around a central chlorine atom. Perchlorate is usually found as the anion component of a salt most often associated with cations such as ammonium, sodium, or potassium. It is typically found in the form of perchloric acid and salts such as ammonium perchlorate, potassium perchlorate, and sodium perchlorate. Perchlorate is a strong oxidizing anion and has recently gained public attention following its discovery in well water and drinking water throughout the United States. Perchlorate is ubiquitous and has been found to be present in soil, vegetation, groundwater and surface water in a number of places. The non-volatile and stable nature of the perchlorate anion has allowed it to be present in drinking water aquifers. It is used as an oxidizer in flares, pyrotechnics, explosives, and numerous other applications, and the recent detection of environmental contamination has primarily been associated with its use in rocket propellants and missile motors. Despite the detection of perchlorate in groundwater in the 1950s, its toxicological properties have only recently been examined in detail [1,2]. The primary toxicity issue is associated with disruption of iodide uptake in the thyroid gland due to its similarity in ionic radius to iodide. Widespread presence of perchlorate in drinking water aquifers and its toxicological properties make perchlorate an emerging chemical of concern.

#### 1.1. Properties of Perchlorate

Commonly used perchlorate compounds are all mostly solid salts at ambient temperature and consist of white or clear crystals (Table 1). Perchlorate salts have low vapour pressure and hence cannot volatilize under ambient conditions. Perchlorate salts are highly soluble in water and dissolved perchlorate anions do not tend to partition from aqueous to gas phase. This soluble, noncomplexing nature makes perchlorate highly mobile in the aqueous environment and does not allow sorption to soils or bioaccumulation. This also implies that perchlorate cannot be treated by precipitation or

sorption, while its nonvolatile nature implies that it cannot be treated by air stripping.

Compound	Formula	Molecular weight (g/mol)	Physical appearance
Ammonium perchlorate	$\text{NH}_4\text{ClO}_4$	117.496	White crystalline solid
Potassium perchlorate	$\text{KClO}_4$	138.55	White crystalline solid
Sodium perchlorate	$\text{NaClO}_4$	120.44	White crystalline solid

[Table 1.](#)

Characteristics and uses of some perchlorate compounds [2,3].

Thermodynamic measurements indicate perchlorate anion as a strong oxidant, however slow kinetics generally limits its reactivity. The lack of reactivity results from high strength of chlorine-oxygen bonds, and the requirement that reduction proceed by removal of an oxygen atom, rather than by direct interaction of a reducing agent with a chlorine atom [2]. This capacity as a strong oxidant and its limited reactivity nature allows perchlorate salts to persist in the environment and they are hard to treat by chemical reduction, while biologically mediated reactions can effectively degrade them.

## 2. Uses of Perchlorate

[Go to:](#)

The most commonly manufactured perchlorate compounds are ammonium, sodium and potassium perchlorate and perchloric acid. Various perchlorate-containing materials are used for specialized applications in chemical industries. Because of its exceptional oxidizing capacity, perchlorate is widely used by the industry, in the manufacture of munitions, explosives and fireworks, a few specific medicinal applications and in small amounts in other high volume consumer products. Perchlorate has been used medically to control hyperthyroid conditions and Graves disease [4,5]. Martino *et al.* [4] studied the effect of potassium perchlorate in patients with hypothyroidism. Short term administration of potassium perchlorate to six patients led to prompt restoration of euthyroidism. Most of the perchlorate-contaminated sites were once associated with rocket manufacturing or munitions use. Use of fireworks, flares and blasting agents has also resulted in environmental contamination [6]. A study conducted on surface water samples following a firework display, found a maximum perchlorate concentration of 44.2  $\mu\text{g/L}$ , while the concentrations decreased towards the background level within 20 to 80 days [7]. Perchlorate may also be formed as a breakdown product of sodium hypochlorite, widely used as a bleach and can be formed incidently in corrosion control applications.

## 3. Environmental Fate and Transport

[Go to:](#)

Perchlorate is highly soluble in water and stable under normal atmospheric conditions. Once a perchlorate compound is released into the environment, most often into water bodies, it gets dissolved in water, and due to its limited ability to adsorb on to surfaces, and/or gets transported through bulk movement of water and mixing processes. When the flow is less or if water becomes stagnant, perchlorate tends to diffuse into clay layers in an aquifer. Perchlorate may be biologically degraded by indigenous microorganisms present in water under anaerobic conditions [8]. Terrestrial and aquatic plants can take up perchlorate and accumulate in various tissues or degrade it in their leaves and branches [9].

## 4. Perchlorate in the Environment

[Go to:](#)

### 4.1. Presence of Naturally Occurring Perchlorate

Most naturally occurring sources of perchlorate appear to be geographically limited to arid environments. These deposits tend to be low in concentration, except for the relatively high natural concentrations found in Chilean caliches and some potash ores [10]. The major Chilean source lies in the Atacama Desert, where the fertilizer is extracted from deposits of nitrate ores or brines [11]. The deposits contain perchlorate ranging from 0.03 to 0.1% or 300 to 1,000 mg/kg in soil [10]. Potash ore is mined and milled in Saskatchewan, Canada (silvite mineral), United States (New Mexico, hankite mineral from California), and Playa crusts from Bolivia [12]. The potassium chloride in these deposits originated in briny sea beds, similar to those of Chilean deposits. The concentrations of perchlorate in these deposits range from 25 to 2,700 mg/kg. Infrequent precipitation in arid environments minimizes the opportunity for perchlorate in nitrate and potash deposits to dissolve and migrate to groundwater. Hence, perchlorate would remain in these deposits in larger quantities than the ones located in regions of higher precipitation.

### 4.2. Theory of Origin of Natural Perchlorate

In addition to occurring in these deposits, a current theory concerning the origin of naturally occurring perchlorate in the environment focusses on natural atmospheric processes. This theory suggests that chloride, possibly in the form of sodium chloride from sea or land based chloride compounds blown in from the atmosphere, reacts with atmospheric ozone to

create perchlorate. This process occurs over much of the earth and is analogous to nitrate formation in the atmosphere [13]. Photolysis of ozone in the presence of sea salt is predicted to form chlorine ( $\text{Cl}_2$ ) and hypochlorous acid ( $\text{HOCl}$ ) in the marine environment [14]. During such reactions, formation of chlorate ( $\text{ClO}_3^-$ ) has been proposed in the stratosphere [15]. The reactions were reexamined on the surface of the sodium chloride particles [16] and studies show that oxygenated products do form even on dry salt surfaces [17]. Chlorate has also been observed to form during ozonation of chloride ( $\text{Cl}^-$ ) containing natural waters. Dissolution of molecular chlorine ( $\text{Cl}_2$ ) in water may result in formation of hypochlorite radicals ( $\text{OCl}^-$ ) [18]. Addition of oxygen to chlorate ( $\text{ClO}_3^-$ ) or fractionation of hypochlorite may result in the formation of perchlorate radical ( $\text{ClO}_4^-$ ) [19]. These reactions are thermodynamically favourable but reasons for not forming appreciable amounts of perchlorate may be kinetic [19]. Given the precursor ingredients and sufficiently energetic conditions, it should be possible to induce formation of perchlorate. A recent theory suggests that in addition, there is a possibility that lightning may play a role in the creation of some atmospherically produced perchlorate [19]. Following atmospheric creation, perchlorate returns to the earth's surface dissolved in precipitation. In arid environments, where the rate of deposition exceeds the rate of dissolution by ongoing precipitation, perchlorate can be incorporated into certain geologic formations [11].

### 4.3. Presence of Perchlorate in Natural Waters

Table 2 presents information on perchlorate release into the environment. Perchlorate has been detected in selected surface waters in the Great Lakes basin in Canada [20]. Of the 55 sites in the Great Lakes basin sampled and analyzed for the presence of perchlorate, it was detected at concentrations near the detection limit in two samples from Hamilton Harbour and six creeks/streams in the Maitland Valley and the Upper Thames River watersheds. The two detections in the Harbour were thought to be related to the fireworks display which had occurred prior to sampling. Presence of low but detectable concentrations of perchlorate, approximately  $1 \mu\text{g/L}$ , has also been confirmed in an unconfined aquifer located in southwestern Ontario, Canada [21]. Most of the perchlorate contamination that has been found in the United States has been associated with military activities or defense contractors. Considerable attention has been focused on Lake Mead and the Colorado River because of the ammonium perchlorate production facilities formerly located in Henderson, Nevada. So far, dense perchlorate occurrence in United States has been found in California, Texas, Massachusetts, Maryland, Oregon and New Jersey. Drinking water in about 26 states in the United States is believed to be contaminated with perchlorate with less than  $12 \mu\text{g/L}$  concentration [2]. A recent study to determine the overall distribution of perchlorate in California drinking water sources found that the mean perchlorate level was  $3.6 \mu\text{g/L}$  and among sources with positive perchlorate results, the mean level was  $12 \mu\text{g/L}$  [22].

Table 2.

Releases of perchlorate compounds [3,26,27].

The occurrence of perchlorate in Japan in the Tone River Basin has been recently detected [23]. Perchlorate was found at high concentrations in the upper Tone River and its tributary, Usui River, and the maximum concentrations were 340 and  $2,300 \mu\text{g/L}$ , respectively. The possible sources of perchlorate in the two areas were attributed to industrial effluents. It was shown that tap waters in the Tone River Basin were also widely contaminated with perchlorate. Widespread contamination of perchlorate in sludge, rice, bottled water and milk collected from different areas in China has been confirmed [24]. The concentration of perchlorate in sewage sludge, rice, bottled drinking water and milk was in the range of  $0.56\text{--}379.9 \mu\text{g/kg}$ ,  $0.16\text{--}4.88 \mu\text{g/kg}$ ,  $0.037\text{--}2.013 \mu\text{g/L}$  and  $0.30\text{--}9.1 \mu\text{g/L}$ , respectively.

### 4.4. Perchlorate in Food Supplies

The primary route of human exposure to perchlorate is via ingestion of perchlorate-contaminated water and food. Ingested perchlorate is readily and completely absorbed from the gastrointestinal tract and excreted rapidly primarily via the urine. Based on available literature, ingestion of perchlorate contaminated drinking water is one of the major exposure routes of concern, although ingestion of contaminated food and human milk are other potential sources of exposure (Table 3). When compared to ingestion, skin absorption and inhalation of perchlorate can be considered negligible exposure pathways.

Table 3.



billion. At MCL of 24 µg/L, the estimated compliance cost drops to \$0.1 billion because of a small number of PWSs contaminated with perchlorate at that level.

## 7. Toxicity and Risk

[Go to:](#)

### 7.1. Thyroid Function

The primary function of thyroid is production of the thyroid hormones, triiodothyronine (T<sub>3</sub>) and thyroxine (T<sub>4</sub>). Iodide is transported into thyroid follicular cells against a concentration gradient by the sodium iodide symporter (NIS) molecule. Once gaining entry to the thyroid follicular cells, iodide is subsequently oxidized to iodine by enzyme thyroglobulin peroxidase (TPO). Iodine is coupled to tyrosine residues on the thyroglobulin (T<sub>g</sub>) molecule [45]. Thyroglobulin is stored within a cavity inside the thyroid follicle in the form of a viscous substance called colloid. In response to signals from the pituitary, T<sub>g</sub> is transported back into follicular cell and is cleaved to yield T<sub>3</sub> and T<sub>4</sub>. Secretion of thyroid hormones is controlled by a well-known feedback mechanism. When serum T<sub>3</sub> and T<sub>4</sub> levels are too low, thyrotropin releasing hormone (TRH) is secreted by the hypothalamus and thyroid stimulating hormone or thyrotropin (TSH) is released by anterior pituitary to promote thyroidal iodide uptake and thyroid hormone synthesis. The subsequent rise in serum T<sub>3</sub> and T<sub>4</sub> levels result in a negative feedback, causing a reduction in circulating levels of TRH and TSH. Measuring serum levels of these hormones represents the standard approach for assessing thyroid function [46].

### 7.2. Perchlorate Toxicity

Perchlorate in drinking water or ingested by other means leads to inhibition of uptake of iodine (as iodide) into thyroid. The perchlorate ion blocks the NIS protein that normally acts as an iodide pump on the surface of the thyroid follicle. This is a competitive inhibition and therefore reversible when exposure to perchlorate ceases. Hence the adverse effect is hypothyroidism. Perchlorate also affects other tissues where NIS is known to be present in the lactating breast epithelium, gastrointestinal tract, placenta, skin and mammary gland. The presence of perchlorate decreases the synthesis of circulating thyroid hormones in the adult and decreases their placental transfer to the fetus in pregnant women. Iodide uptake inhibition is considered the mode of action for perchlorate. The mode of action agreed to by toxicologist and risk assessors identifies that inhibition of iodine uptake by perchlorate in thyroid will decrease serum T<sub>3</sub> and T<sub>4</sub> and increase TSH levels resulting in thyroid hypertrophy or hyperplasia, which ultimately would lead to hypothyroidism. The mode of action also identifies that perchlorate has a threshold for effects and that the degree of effects are dependant on dosage. The effects of perchlorate on thyroid function require large doses. Animal and clinical studies conducted, were mostly based on this mode of action for perchlorate.

### 7.3. Animal Toxicity Studies

To determine the health effects of perchlorate in humans, fairly extensive animal toxicity studies have been conducted. The amounts of perchlorate used in these animal studies are huge, relative to potential human exposure. In a number of sub-chronic perchlorate studies, increased thyroid follicular cell hypertrophy and hyperplasia were observed in some of the treated animals [46–48]. These data indicate that oral administration of perchlorate induces hyperplasia in the thyroid of rodents and if the exposures are lengthened, some of the lesions might progress to cancerous thyroid tumours [49,50]. Thyroid follicular tumours were observed in the F<sub>1</sub> generation during a two-generation study in Sprague–Dawley rats [47]. While these studies suggest the potential for thyroid carcinogenicity associated with disruption of thyroid physiology, evidence of an early onset of tumourigenesis in the offspring of treated dams (female ancestors) requires attention because of the suggestion for potential adverse effects in the fetus. The genotoxicity studies have established that ammonium perchlorate does not have significant mutagenic activity [51,52].

There are no chronic exposure studies available in animals. There is clear evidence that perchlorate exposure at elevated doses does produce tumours, and it should be noted that perchlorate is carcinogenic in laboratory animals. The mode of action of perchlorate is established to be operative in laboratory animals and is considered relevant to humans. Rats and human thyroid work similarly and rat toxicity studies are valuable tools for qualitative information regarding the human thyroid. However, there are physiological differences between the human and rat pituitary thyroid axis, which make rats inappropriate for quantifying predicted changes in humans for risk assessment purposes. In humans, the pool of thyroxine-binding-globular thyroid hormone acts as a stable reserve to be called upon when required to supplement free thyroxine. Rats, which have no high affinity carrier do not have this reserve capacity, and compensate for this by increased TSH production and a higher rate of turnover of thyroid hormones [47].

#### 7.4. Human Toxicity Studies

Humans who receive administered doses of perchlorate demonstrate rapid absorption that results in the appearance of peak serum concentrations within approximately three hours. The half-life of perchlorate in human serum is about 6 to 8 hours. Perchlorate is rapidly eliminated unchanged from the body via urinary excretion. Various studies have described the effects of perchlorate on thyroid function in healthy adult humans. In a study by Brabant *et al.* [53] five healthy men were each given daily doses of 13 mg/kg of potassium perchlorate. No differences were noted for T<sub>4</sub> or T<sub>3</sub> at the termination of each four-week exposure period. Brabant *et al.* [53] concluded that there was a change in iodine uptake during the perchlorate treatment. In a second study, Lawrence *et al.* [54] recruited 9 healthy men and had them ingest 10 mg/L of potassium perchlorate daily for fourteen days. There were no notable changes in T<sub>4</sub>, T<sub>3</sub> or serum free TSH during the period of perchlorate exposure. Lawrence *et al.* [55] expanded their earlier perchlorate exposure study by changing the ingestion exposure for eight men to 3 mg of potassium perchlorate daily, again for fourteen days. During perchlorate treatment, radioiodide uptake was found to decrease in its value. Again there were no notable changes in T<sub>4</sub>, T<sub>3</sub> or serum free TSH. Braverman *et al.* [56] observed that a six month exposure to perchlorate by thirteen healthy volunteers at doses up to 3 mg/d had no effect on thyroid function. The most comprehensive study of the kinetics of uptake of iodide and inhibition by perchlorate in humans has been described by Greer *et al.* [41]. In this study, 21 healthy women and 16 healthy men were given potassium perchlorate at several doses ranging from 0.007 mg/kg-day to 0.5 mg/kg-day for fourteen days. Greer *et al.* [41] identified the lowest exposure level (0.007 mg/kg-day) as a no observed adverse effect level (NOAEL) for exposure to perchlorate. There was no evidence for changes in free serum T<sub>4</sub>, T<sub>3</sub> or TSH in any of the participants at doses less than 0.5 mg/kg-day during the study. Blount *et al.* [57] assessed the effect of low-level exposure to perchlorate on thyroid function in 2,299 men and women in the United States during 2001–2002. The study evaluated the potential relationship between urinary levels of perchlorate and serum levels of TSH and T<sub>4</sub>. The results indicated a small increase in serum TSH and small decrease in serum T<sub>4</sub> only in women with urine concentrations < 100 µg/L. Perchlorate was found to have no effect on TSH and T<sub>4</sub> in men. In contrast, Pearce *et al.* [58] found no change in thyroid function in first trimester pregnant women in Europe whose iodine intake was < 100 µg/L. Another study used individuals who received occupational exposures to perchlorate [59]. This study compared workers and members from the surrounding community (non-exposed) at an ammonium perchlorate manufacturing facility. A total of twenty-nine perchlorate workers and twelve community controls participated in the study. Exposure doses were predicted for each shift from assumed 8-hour half-life of perchlorate. It was reported that worker radioiodide uptake values measured during exposure periods were significantly lower than their baseline pre-exposure values. Thyroid hormone levels showed no differences in TSH but higher concentrations of T<sub>3</sub> or T<sub>4</sub> for post-shift exposure was observed.

#### 7.5. Ecotoxicity Studies

In contrast to the abundance of human health data, few ecological data exist on the potential exposure and effects of perchlorate in the environment [60]. There are especially few data regarding the presence and effects of perchlorate on invertebrates, such as earthworms. The toxicity of sodium perchlorate to *Eisenia fetida* was evaluated in sub-chronic tests in artificial soil. The sub-chronic (21 day sand tests) EC<sub>50</sub> for sodium perchlorate to *E. fetida* reproduction (cocoon production) was 1.3 µg/g. The sub-chronic (28 day artificial soil tests) EC<sub>50</sub> for sodium perchlorate to *E. fetida* reproduction (cocoon production) was 350 µg/g [61]. To investigate chronic toxicity in fishes, mosquitofish (*Gambusia holbrooki*) adults and fry were exposed to aqueous sodium perchlorate at 1, 10, and 100 mg/L, and growth and reproductive performance were determined [62]. Five-day acute toxicity tests were also performed. The LC<sub>50</sub> of sodium perchlorate was 404 mg/L. Growth was enhanced at 1 mg/L but inhibited at 10 mg/L. These results suggest that, at environmentally relevant concentrations, perchlorate does not induce acutely toxic effects but may have mild stimulatory or hormetic effects on fitness parameters in this species. Adult female and male zebrafish were paired in water containing ammonium perchlorate at 0, 18, and 677 mg/L for up to 8 weeks [63]. Perchlorate does not accumulate in whole fish. Thyroid follicle hypertrophy and angiogenesis were observed for the 677 mg/L treatment group for 4 weeks, which may be due to extra thyroidal toxicity. More pronounced effects including hypertrophy, angiogenesis, hyperplasia, and colloid depletion were observed for the 18 mg/L treatment group for 8 weeks. Goleman *et al.* [64] evaluated the effects of ammonium perchlorate at environmentally relevant concentrations on development and metamorphosis in amphibian, *Xenopus Laevis* after 70-day exposure. No concentration-dependent developmental abnormalities were observed below the 70-day LC<sub>50</sub> (223 mg/L). Delays in metamorphosis in *Xenopus laevis* at concentrations ranging from 0.005 to 0.1 mg/L and inhibition of metamorphosis at 0.147 mg/L have been reported. It was concluded that

perchlorate may pose a threat to normal development and growth in natural amphibian populations. Birds whose migratory routes transect contaminated watersheds or those whose movement is restricted may be exposed to perchlorate. Thyroid glandular T4 content was significantly decreased in 21-day bobwhite quail embryos from eggs with a concentration of  $\geq 0.5 \mu\text{g/g}$  ammonium perchlorate and in hatchlings from eggs with  $\geq 50 \text{ mg/L}$  ammonium perchlorate [65]. However, egg concentrations of up to  $300 \mu\text{g/g}$  did not result in hypothyroidism in embryos or hatchlings. Perchlorate sensitivity also appears to be species specific in birds [66]. In areas where perchlorate-contaminated groundwater is used for irrigation and/or a source of drinking water, livestock may consume considerable amounts of perchlorate on a daily basis. Perchlorate exposure in cattle was evaluated by monitoring heifer calves on a site with access to perchlorate contaminated water ( $25 \mu\text{g/L}$ ) for 14 weeks. Perchlorate was detected in blood twice ( $15 \mu\text{g/L}$  and  $22 \mu\text{g/L}$ ) in one of the heifer calves at week 4 and week 6, respectively [67].

## 8. Natural Biodegradation

Go to:

Aslander [68] was the first to observe microbial degradation of chlorate in soils. Later, it was shown that bacteria capable of perchlorate and chlorate respiration exist in nature [69,70]. van Ginkel *et al.* [70] demonstrated qualitatively that chlorate respiring microorganisms were present in a variety of environments such as wastewaters, rivers, sediments and soils. These studies suggest that chlorate and perchlorate are biodegradable under certain conditions. Several bacterial isolates have been shown to be capable of completely reducing perchlorate to chloride for cell respiration. The details of perchlorate reducing strains and the reduction process are discussed in detail under the treatment technologies section. Although, perchlorate respiring bacteria are widely distributed in nature, perchlorates still persist in the environment. Intrinsic remediation of perchlorate contaminated site is possible if perchlorate respiring microorganisms exist naturally in the system and can compete for existing sources of organic matter. Otherwise, perchlorate reduction must be enhanced by addition of nutrients to the contaminated sites. Wu *et al.* [71] observed that the presence of perchlorate in the soil tested appears to have enriched the number of perchlorate and chlorate respiring microorganisms in that soil. Perchlorate respiring microorganisms were found to be present in sufficient numbers to achieve complete perchlorate degradation using lactate in soil microcosms in only 7 days.

## 9. Treatment Technologies

Go to:

Widely used treatment technologies used for removing perchlorate from groundwater include, ion exchange, use of bioreactors and in-situ bioremediation. Groundwater contamination of perchlorate is of concern than contamination in soil due to the highly soluble nature of perchlorate in water. Hence the following discussions apply to groundwater and drinking water treatment.

### 9.1. Ion Exchange (IX)

Ion exchange (IX) is an *ex situ* technology used to remove perchlorate from groundwater and surface water. The most commonly used ion exchange media are synthetic, strongly basic, anion exchange resins. Extensive research has focused on developing IX resins that would selectively remove perchlorate when the water contains other competing anions, such as chloride, sulphate, nitrate and bicarbonate, at higher concentrations. Strong base anion-exchange resins have proven to be very effective in removing perchlorate from water [72]. The functional groups in perchlorate resins have positive charges, which are initially dosed with anions such as chloride ion that attaches itself to these positively charged functional groups. The perchlorate anion in the influent water is also attracted to the positively charged functional groups, due to its stronger attractive force and replaces the chloride ion [73]. Thus ion exchange removes the perchlorate ion from the aqueous phase by exchange with the chloride ion, populating the treated water with chloride ions. The regeneration solutions used to regenerate the exhausted resin include 12% brine solution of sodium chloride, which can recapture approximately 7% of the perchlorate from the resin, and ferric chloride in solution with hydrochloric acid, which can recapture close to 100% of the perchlorate from the resin [74]. Many groundwater and drinking water systems incorporate ion-exchange to treat water containing 10 to  $10,000 \mu\text{g/L}$  perchlorate (Table 5). There are fifteen full scale applications of ion exchange so far to remove perchlorate.

Name of technology	Description	Performance	Reference
Ground water	A cation-exchange resin	100%	[72]
Drinking Water	A mixed bed, non-regenerable resin	100%	[74]

Table 5.

A summary of performance of ion exchange used for treatment of perchlorate.

## 9.2. Bioreactors

A bioreactor is an *ex situ* biological treatment system for degrading perchlorate from contaminated groundwater and surface water using microorganisms. Certain bacteria can reduce perchlorate in soil and water under certain anaerobic conditions. The microorganisms most often used in perchlorate treatment are facultative anaerobes that use electron acceptors such as nitrate, sulphate and perchlorate to reduce organic compounds. In addition to the electron sink, these organisms require carbon sources such as ethanol, methanol, or acetic acid for growth. Groundwater naturally contains some perchlorate reducing bacteria [8] and under reducing conditions such as in a bioreactor, population of such bacterial species will increase over time as the microbes select perchlorate as the dominant electron acceptor. Addition of carbon sources induces the microorganisms in water to extract electrons from the carbon source and deliver them to perchlorate. This electron transfer induces biotransformation of perchlorate ion to chlorate and then chloride [2]. Recent research shows that perchlorate reduction involves a three step process catalyzed by two enzymes. A perchlorate reductase enzyme catalyzes reduction of perchlorate to chlorate and then to chlorite. A chlorite dismutase enzyme causes further breakdown of chlorite to chloride and oxygen [77,78]. The rate limiting step is believed to be perchlorate reduction to chlorate [79]. Perchlorate reducing strains reported in the literature include *Vibrio dechloraticans* Cuznesnove B-1168 [80], *Wolinella succinogens* HAP-1 [81], and isolates GR-1 [82], perclace [61], *Dichlorosoma* sp. KJ [83] and *Dechloromonas agitata* strain CKB [84]. Several different types of bioreactors have been proposed to remove perchlorate from water using mixed and pure cultures of bacteria [85]. Table 6 gives a summary of various bioreactors used for treating perchlorate containing water. Fluidized bed reactors (FBR) and packed bed reactors (PBR) are two types of bioreactors widely used for removal of perchlorate. FBRs are generally made up of suspended sand or granular activated carbon media to support microbial activity and biomass growth. The biomass on the packing is kept uniform in the reactor by mixing, and biomass thickness is controlled by shear and particle collisions. Packed bed reactors are made up of static sand or plastic media to support the growth of microbes. Perchlorate has been degraded in saturated-flow packed bed reactors, air lift reactors and unsaturated flow packed bed reactor [87,88]. The overall rate of perchlorate degradation in these reactors is first order with respect to log mean perchlorate concentration in the reactor [85]. The highest perchlorate removal rate in a fixed bed reactor was obtained for a sand packed reactor using *Dechlorosoma* sp. KJ [83]. Studies have been conducted on the use of hydrogen gas reactors, which use continuous hydrogen gas transfer to the biofilm with unsaturated flow conditions for autotrophic perchlorate reduction [88,89]. Hydrogen oxidizing bacterial isolate, designated as JM has been used in these studies.

Type of reactor	Description	Reference
Batch	Batch reactor with 100 ml of water containing 10 mg of perchlorate and 10 mg of carbon source.	[80]
Batch	Batch reactor with 100 ml of water containing 10 mg of perchlorate and 10 mg of carbon source.	[81]
Batch	Batch reactor with 100 ml of water containing 10 mg of perchlorate and 10 mg of carbon source.	[82]
Batch	Batch reactor with 100 ml of water containing 10 mg of perchlorate and 10 mg of carbon source.	[61]
Batch	Batch reactor with 100 ml of water containing 10 mg of perchlorate and 10 mg of carbon source.	[83]
Batch	Batch reactor with 100 ml of water containing 10 mg of perchlorate and 10 mg of carbon source.	[84]
Batch	Batch reactor with 100 ml of water containing 10 mg of perchlorate and 10 mg of carbon source.	[85]
Batch	Batch reactor with 100 ml of water containing 10 mg of perchlorate and 10 mg of carbon source.	[87]
Batch	Batch reactor with 100 ml of water containing 10 mg of perchlorate and 10 mg of carbon source.	[88]
Batch	Batch reactor with 100 ml of water containing 10 mg of perchlorate and 10 mg of carbon source.	[89]

**Table 6.**

Summary of Bioreactor studies for perchlorate treatment.

## 9.3. In situ Biodegradation

*In situ* biodegradation (ISB) can be used for source reduction, dissolved phase contaminant reduction or as a biological barrier to contain a plume [93]. ISB involves delivery of nutrients such as ethanol, lactate, acetate, citrate, sugars and edible oils to the subsurface to promote the biodegradation of perchlorate by the indigenous bacteria. The most important consideration in ISB is the ability to transmit and mix liquids in the subsurface [93]. Major system components include extraction wells, conveyance piping, electron donor dosing station, electron donor delivery/recharge wells, barrier wall width and depth, barrier components and performance monitoring wells. Electron donors can be delivered to the aquifer through a variety of active (recirculation) or passive (injection or trenching/barrier) methods [2]. Permeable reactive barriers (PRB) can be used to deliver the amendments to the contaminated groundwater. PRBs are installed as permanent, semi-permanent, or temporary units across the flow path of the contaminant plume. Contaminants in the groundwater that flow through a PRB are degraded chemically or biologically. This biobarrier approach can use engineered trenches or barriers containing solid phase [94] or viscous substrates [95] placed across the flow path through direct push injections. Examples of reactive materials used include edible oils, pecan shells, cotton seed, chitin, limestone, and other composting materials. Another approach to deliver electron donors include mobile amendment systems, which are characterized by injecting less viscous water soluble substrates into the up-flow of the contaminant plume, allowing the substrate to move down treating the groundwater as it moves. Active treatment can use groundwater extraction and reinjection system to distribute substrate in the subsurface [96] or horizontal-flow treatment wells to mix and distribute electron donors in groundwater [76]. Table 7 provides a list of ISB technology used for treating perchlorate contaminated water.

Table 7.

Summary of in situ bioremediation for treating perchlorate.

#### 9.4. Other Processes

1. Adsorption: Adsorption involves the use of adsorbent media such as granular activated carbon (GAC) or activated alumina to remove perchlorate. For sites that have groundwater containing contaminants other than perchlorate, previous experience shows the potential for the use of treatment trains consisting of standard GAC/ion exchange resins or tailored GAC [97]. Cationic surfactant-tailored activated carbon has been studied to remove perchlorate in a laboratory scale column test; perchlorate was removed to below detection levels and increased the breakthrough time up to 30 times longer than standard activated carbon [98].
2. Membrane filtration: Membrane filtration treatment includes reverse osmosis (RO), nanofiltration (NF), ultrafiltration (UF) and electrodialysis (ED). Three main types of semi-permeable membranes have been evaluated for the treatment of perchlorate in groundwater which includes high pressure RO membrane, nanofiltration membrane and low pressure RO membrane. High pressure RO membranes (> 150 psi) have been observed to remove about 99.9% of perchlorate [99]. NF membranes have a nominal pore size of approximately 0.001 micron and require an operating pressure in the range of 80 to 150 psi. NF membranes have been found to allow particles such as salts to pass through them. UF membranes have a pore size of approximately 0.002 to 0.1 microns and operate at a lower pressure (30 to 100 psi). Tight UF membranes, having smaller pores than UF but larger than NF, have been found to be ineffective in treating perchlorate [2]. Liang *et al.* [100] conducted pilot scale studies to evaluate the performance of perchlorate removal by production sized RO, NF and UF membranes. RO and NF membranes were observed to consistently remove more than 86.7% of perchlorate while tight UF membranes proved to be ineffective. Addition of cationic surfactant to UF membrane did not prove to be effective; this resulted in plugging the membrane.
3. Electrochemical reduction: Electrochemical reduction of perchlorate anion has been studied in a cell with a nickel working electrode and a platinum counter electrode in concentrated solutions of hypochlorous acid [102]. Lee and Kramer [103] examined titanium metal as a chemical reductant to remove perchlorate in water. The activation of titanium was achieved by eliminating the localized surface oxide film using electrochemically induced pitting corrosion. It was observed that a higher current increases the activity of pitting corrosion. This dissolves a higher concentration of transitory titanium metal ions in the vicinity of the pits, which results in a higher rate of perchlorate reduction. The surface of the bare Ti(0) inside the pits induces further electrochemical reactions and causes faster rate of chloride oxidation to chlorine by increasing the current.
4. Phytoremediation: Phytoremediation is an in situ mechanism by which vegetation is used to treat perchlorate. The most successful plants in perchlorate degradation include French tarragon, cottonwood, and willow. Under hydroponic conditions willows were observed to degrade perchlorate from 10,000 µg/L to below detection in about 53 days [9]. Other plants that have been successfully tested include salt cedar trees, bulrushes, cattails and sedges. Perchlorate concentrations have decreased from 20 µg/L to less than 4 µg/L using constructed wetlands [2].
5. Use of iron particles: Stabilized elemental iron nanoparticles have been studied to remove perchlorate [104]. The results showed that temperature played a critical role in perchlorate degradation process. Approximately 100% of perchlorate was transformed to chloride during the reaction without any detectable intermediate products. Kinetics of perchlorate reduction by elemental iron was examined at elevated temperatures using microwave heating and conventional block heating [105]. Results from microwave heating study showed that 98% of perchlorate was removed in one hour at 200 deg C. Similar results observed with block heater indicated that enhancement in the rate of perchlorate removal by elemental iron was mostly due to heat energy at a higher temperature. Gurol and Kim [106] investigated chemical reduction of perchlorate using iron and iron oxide under various conditions. The main result of their investigation was that the addition of metallic iron (100 g/L) combined with exposing the solutions to ultraviolet light at wavelengths < 185 nm achieved 77% reduction in perchlorate.
6. Catalytic reactors: A preliminary study using a catalyst that efficiently uses hydrogen gas to reduce perchlorate completely to chloride has been reported [107]. The catalyst has been prepared by the addition of methylthioxorhenium to a combination of 5% Pd-carbon powder. Initial batch studies have indicated complete

reduction of perchlorate to chloride.

Studies such as thermal destruction, capacitative deionization have also been reported in literature as effective treatment systems for perchlorate.

## 10. Conclusions

[Go to:](#)

Perchlorate has been found to be widespread in drinking water sources in the United States and a few other countries. Recent studies have confirmed the presence of perchlorate in Canadian surface waters. Though, known adverse effect of perchlorate exposure on humans is hypothyroidism, detailed animal toxicity studies conducted on two generation of rats have shown to have potential adverse effects on their fetuses. Perchlorate has been found in cow's milk and mother's milk and hence perchlorate ingestion by infants through maternal intake is high. Therefore, perchlorate can be considered as a contaminant of concern in the near future in Canada.

The most common perchlorate treatments currently in use employ ex situ technologies. Two treatment technologies currently dominate the field of perchlorate treatment: ion exchange and biological treatment. Research demonstrated that bioreactors, mostly fluidized bed reactors operated under anaerobic conditions, could effectively reduce high concentrations of perchlorate. Selective ion exchange resins of various types have been developed to treat lower concentrations of perchlorate. Although bioreactors have been proven to be effective, the best applicable technology for removal of perchlorate remains to be ion exchange. Ion exchange resins are used for treatment of drinking water supplies and groundwater containing 10 to 100,000 µg/L perchlorate. Research is presently directed on developing technologies that could remove perchlorate to less than 10 µg/L from water sources. Such recent technologies include electrochemical reduction, catalytic reduction and use of metallic nanoparticles.

## References

[Go to:](#)

1. Lamm S, Doemland M. Has perchlorate in drinking water increased the rate of congenital hypothyroidism? *J. Occup. Environ. Med.* 1999;41:409–411. [[PubMed](#)]
2. Sellers K, Alsop W, Clough S, Hoyt M, Pugh B, Robb J, Weeks K. *Perchlorate: Environmental problems and solutions.* CRC Press; Boca Raton, FL, USA: 2006. p. 177.
3. Motzer W. Perchlorate: problems, detection and solutions. *Environ. Forensics.* 2001;2:301–311.
4. Martino E, Mariotti S, Aghini-Lombardi F, Lenziardi M, Morabito S, Baschieri L, Pinchera A, Braverman L, Safran M. Short term administration of potassium perchlorate restores euthyroidism in amiodarone iodine-induced hypothyroidism. *J. Clin. Endocrinol. Metab.* 1986;63:1233–1236. [[PubMed](#)]
5. Wenzel KW, Lente JR. Similar effects of thionamide drugs and perchlorate on thyroid stimulating immunoglobulins in Graves' disease: evidence against an immunosuppressive action of thionamide drugs. *J. Clin. Endocrinol. Metab.* 1984;58:62–69. [[PubMed](#)]
6. Cox C. Statistical issues for animal studies of developmental neurotoxicity. In: Weiss B, O'Donoghue JL, editors. *Neurobehavioral toxicity: analysis and interpretation.* Raven Press; New York, NY, USA: 1994. pp. 93–101.
7. Wilkin RT, Fine DD, Burnett NG. Perchlorate Behavior in a Municipal Lake Following Fireworks Displays. *Environ. Sci. Technol.* 2007;41:3966–3971. [[PubMed](#)]
8. Coates JD, Michaelidou U, Bruce RA, O'Connor SM, Crespi JN, Achenbach LA. The ubiquity and diversity of dissimilatory (per)chlorate reduction. In: Urbanski T, editor. *Perchlorate in Environment.* Kluwer Academic/Plenum Publishers; New York, NY, USA: 1999. pp. 257–270.
9. Nzungu VA, Wang C, Harvey G. Plant mediated transformation of perchlorate into chloride. *Environ. Sci. Technol.* 1999;33:1470.
10. Duncan PB, Morrison RD, Vavricka E. Forensic identification of anthropogenic and naturally occurring sources of perchlorate. *Environ. Forensics.* 2005;6:205–215.
11. Urbanski ET, Collette TW, Robarge WP, Hall WL, Skillen JM, Kane PF. *Survey of Fertilizers and Related Materials for Perchlorate (ClO<sub>4</sub><sup>-</sup>)* Environmental Protection Agency; Cincinnati, OH, USA: 2001. EPA/600/R-01/047.

12. Orris GJ. Perchlorate in natural minerals and materials. USGS Quarterly Report (April). 2004. Available online: [http://www.itrcweb.org/Documents/Perch\\_private/USGS\\_Minerals.doc](http://www.itrcweb.org/Documents/Perch_private/USGS_Minerals.doc) (accessed on August 20, 2007).
13. Walvoord MA, Phillips FM, Stonestrom DA, Evans RD, Hartsough PC, Newman BD, Striegl RG. A reservoir of nitrate beneath desert soils. *Science*. 2003;302:1021–1024. [[PubMed](#)]
14. Oum KW, Lakin MJ, DeHaan DO, Brauers T, Finlayson-Pitts BJ. Formation of molecular chlorine from the photolysis of ozone and aqueous sea-salt particles. *Science*. 1998;279:74–77. [[PubMed](#)]
15. Simonaitis R, Heicklen J. Perchloric acid: A possible sink for stratospheric chlorine. *Planet. Space Sci*. 1975;23:1749–1752.
16. Knipping EM, Dabdub D. Modeling Cl<sub>2</sub> formation from aqueous NaCl particles: Evidence for interfacial reactions and importance of Cl<sub>2</sub> decomposition in alkaline solution. *J. Geophys. Res.* 2002;107:4360.
17. Newberg JT, Hemminger JC. X-ray photoelectron spectroscopy analysis of ozone reacting with halide salt surfaces. Proceedings of 228th ACS National Meeting; American Chemical Society; Philadelphia, PA, USA. 22–26; Aug. 2004.
18. Siddiqui MS. Chlorine-ozone interactions: Formation of chlorate. *Water Res.* 1996;30:2160–2170.
19. Dasgupta PK, Nartubekabgim PJ, Jackson WA, Anderson TA, Tian K, Tock RW, Rajagopalan S. The origin of naturally occurring perchlorate: The role of atmospheric processes. *Environ. Sci. Technol.* 2005;39:1569–1575. [[PubMed](#)]
20. Backus SM, Klawuun P, Brown S, D'sa I, Sharp S, Surette C, Williams DJ. Determination of perchlorate in selected surface waters in the Great Lakes Basin by HPLC/MS/MS. *Chemosphere*. 2005;61:834–843. [[PubMed](#)]
21. Robertson WD, Ptacek CJ, Brown SJ. Geochemical and hydrogeological impacts of a wood particle barrier treating nitrate and perchlorate in groundwater. *Ground Water Monit. Remediat.* 2007;27:85–95.
22. Kimbrough DE, Parekh P. Occurrence and co-occurrence of perchlorate and nitrate in California drinking water sources. *J. Amer. Water Work Assn.* 2007;99:126–132.
23. Kosaka K, Asami M, Matsuoka Y, Kamoshita M, Kunikane S. Occurrence of perchlorate in drinking water sources of metropolitan area in Japan. *Water Res.* 2007;41:3474–3482. [[PubMed](#)]
24. Shi Y, Zhang P, Wang Y, Shi J, Cai Y, Mou S, Jiang G. Perchlorate in sewage sludge, rice, bottled water and milk collected from different areas in China. *Environ. Int.* 2007;33:955–962. [[PubMed](#)]
25. Silva MA. Safety Flares Threaten Water Quality with Perchlorate. Santa Clara Valley Water District; San Jose, CA, USA: 2003.
26. Hjeresen DS, Rae B, Beers M, Saladen CL, Nylander A, Barr A, Pope J, Dziewinski J, Scott R, Holcomb D, Hollis L, Dale L, Williams B, Strietelmeier B, Carlson R, Alexander P, Worland S, Hanson J, Stine M, Hiskey J, Archuleta S, Kinkead A, Sherrard P, Longmire M, Witkowski, Gard M. Los Alamos National Laboratory Perchlorate Issues Update. Los Alamos National Laboratory; Los Alamos, NM, USA: 2003. LA-CP-03-0441.
27. Explanatory data on perchlorate in food. Office of Plant and Dairy Foods, USFDA; Maryland, USA: 2004.
28. Ellington JJ, Wolfe NL, Garrison AW, Evans JJ, Avants J, Teng Q. Determination of perchlorate in tobacco plants and tobacco products. *Environ. Sci. Technol.* 2001;35:3213–3218. [[PubMed](#)]
29. Yu L, Canas JE, Cobb GP, Jackson WA, Anderson TA. Uptake of perchlorate in terrestrial plants. *Ecotoxicol. Environ. Safety.* 2004;58:44–49. [[PubMed](#)]
30. Jackson WA, Joseph P, Laxman P, Tan K, Smith PN, Yu L, Anderson TA. Perchlorate accumulation in forage and edible vegetation. *J. Agr. Food Chem.* 2005;53:369–373. [[PubMed](#)]
31. Sanchez CA, Crump KS, Krieger RI, Khandaker NR, Gibbs JP. Perchlorate and nitrate in leafy vegetables of North America. *Environ. Sci. Technol.* 2005;39:9391–9397. [[PubMed](#)]
32. Kirk AB, Smith EE, Tian K, Anderson TA, Dasgupta PK. Perchlorate in milk. *Environ. Sci. Technol.*

2003;37:4979–4981. [[PubMed](#)]

33. Kirk AB, Martinelango PK, Tian K, Dutta A, Smith EE, Dasgupta PK. Perchlorate and iodide in dairy and breast milk. *Environ. Sci. Technol.* 2005;39:2011–2017. [[PubMed](#)]

34. Pearce EN, Leung AM, Blount BC, Bazrafshan HR, He X, Pino S, Valentin-Blasini L, Braverman LE. Breast milk iodine and perchlorate concentrations in lactating Boston-area women. *J. Clin. Endocrinol. Metab.* 2007;92:1673–1677. [[PubMed](#)]

35. Hautman DP, Munch DJ, Eaton AD, Haghani AW. Method 3140--Determination of Perchlorate in Drinking Water Using Ion Chromatography, Revision 1; US Environmental Protection Agency; Cincinnati, OH, USA: 1999. Doc No EPA/815/B-99/003.

36. Method 9058--Determination of perchlorate using ion chromatography with chemical suppression conductivity detection. USEPA; Washington, DC, USA: 2000.

37. Coleman M, Coates J. Chlorine isotopic characterization of perchlorate. Perchlorate in California's Groundwater. Proceedings of The Eleventh Symposium in GRA's Series on Groundwater Contaminants; Groundwater Resources Association; Glendale, CA, USA. 4; Aug, 2004.

38. Sturchio N, Hatzinger P, Arkins M, Suh C, Heraty L. Chlorine isotope fractionation during microbial reduction of perchlorate. *Environ. Sci. Technol.* 2003;37:3859–3863. [[PubMed](#)]

39. Health Canada Perchlorate and human health. 2005. Available online: [http://www.hc-sc.gc.ca/ewh-semt/water-eau/drink-potab/perchlorate\\_e.html](http://www.hc-sc.gc.ca/ewh-semt/water-eau/drink-potab/perchlorate_e.html) (accessed July 30, 2007).

40. NRC . Health implications of perchlorate ingestion. National Academies Press; Washington, DC, USA: 2005. Health implications of perchlorate ingestion Committee to Assess the Health Implications of Perchlorate Ingestion, National Research Council.

41. Greer MA, Goodman G, Pleus RC, Greer SE. Health effects assessment for environmental perchlorate contamination: The dose response for inhibition of thyroidal radioiodine uptake in humans. *Environ. Health Perspect.* 2002;110:927–937. [[PMC free article](#)] [[PubMed](#)]

42. USEPA . Interim drinking water health advisory for perchlorate 2008. USEPA; Washington, DC, USA: EPA 822-R-08-025.

43. Russel CG, Robertson A, Chowdhury Z, McGuire MJ. National cost implications of a perchlorate regulation. *J. Amer. Water Work Assn.* 2009;101:54–67.

44. USEPA . State perchlorate advisory levels as of 04/20/05. USEPA; Washington, DC, USA: 2005. Available online: <http://www.epa.gov/fedfac/pdf/stateadvisorylevels.pdf> (accessed 2009).

45. Argus Research Laboratories, Inc. Oral (drinking water) developmental toxicity study of ammonium perchlorate in rabbits. Final Pilot Report; Argus Research Laboratories, Inc.; Warminster, PA, USA: 1998.

46. Lewandowski TA, Seeley MR, Beck BD. Interspecies differences in susceptibility to perturbation of thyroid homeostasis: a case study with perchlorate. *Regul. Toxicol. Pharmacol.* 2004;39:348–362. [[PubMed](#)]

47. Keil D, Waren A, Bullard-Dillard K, Jenny M, Eudaly J. Effects of ammonium perchlorate on immunological, hematological, and thyroid parameters. Department of Medical Laboratory Sciences, Medical University of South Carolina; Charleston, South Carolina, USA: 1998.

48. Caldwell DJ, King JH, Kinkead ER, Jr, Wolfe RE, Narayanan L, Mattie DR. Results of a fourteen day oral dosing toxicity study of ammonium perchlorate Proceedings of the JANNAF Safety and Environmental Protection Subcommittee Meeting, Florida, USA December 1995. Vol 1

49. Pajer Z, Kalisnik M. The effect of sodium perchlorate and ionizing irradiation on the thyroid parenchymal and pituitary thyrotropic cells. *Oncology.* 1991;48:317–320. [[PubMed](#)]

50. Kessler FJ, Kruskemper HJ. Experimentelle Schilddrusentumoren durch mehrjährige Zufuhr von Kaliumperchlorat.

[Experimental thyroid tumors caused by long-term administration of potassium perchlorate.] *Klin Wochenschr.* 1966;44:1154–1156. [[PubMed](#)]

51. Zeigler E. Salmonella mutagenicity testing of ammonium perchlorate. US Department of Health and Human Services, National Institute of Environmental Health Sciences; North Carolina, USA: 1998.

52. ManTech Environmental Toxicology, Inc. Genotoxicity assays for ammonium perchlorate. Final report; ManTech Environmental Toxicology, Inc.; Cincinnati, OH, USA: 1998. (cited in USEPA, 1998).

53. Brabant G, Bergmann P, Kirsch CM, Kohrle J, Hesch RD, von zur Muhlen A. Early adaptation of thyrotropin and thyroglobulin secretion to experimentally decreased iodine supply in man. *Metabolism.* 1992;41:1093–1096. [[PubMed](#)]

54. Lawrence JE, Lamm SH, Pino S, Richman K, Braverman LE. The effect of shortterm low-dose perchlorate on various aspects of thyroid function. *Thyroid.* 2000;10:659–663. [[PubMed](#)]

55. Lawrence J, Lamm S, Braverman LE. Low dose perchlorate (3 mg daily) and thyroid function. *Thyroid.* 2001;11:295. [[PubMed](#)]

56. Braverman LE, Pearce EN, He X, Pino S, Seeley M, Beck B, Magnani B, Blount BC, Firek A. Effects of six months of daily low-dose perchlorate exposure on thyroid function in healthy volunteers. *J. Clin. Endocrinol. Metab.* 2006;91:2721–2724. [[PubMed](#)]

57. Blount BC, Pirkle JL, Osterloh JD, Valentin-Blasini L, Caldwell KL. Urinary perchlorate and thyroid hormone levels in adolescent and adult men and women living in the United States. *Environ. Health Perspect.* 2006;114:1865–1871. [[PMC free article](#)] [[PubMed](#)]

58. Pearce EN, Lazarus JH, Smyth PPA, He X, Dall'Amico D, Parkes AB, Burns R, Smith DF, Maina A, Leung AM, Braverman LE. Thyroid function is not affected by environmental perchlorate exposure in first trimester pregnant women. *Thyroid.* 2007;17:S-133. (abstract only).

59. Braverman LE, He X, Pino S, Cross M, Magnani B, Lamm SH, Kruse MB, Engel A, Crump KS, Gibbs JP. The effect of perchlorate, thiocyanate, and nitrate on thyroid function in workers exposed to perchlorate long-term. *J. Clin. Endocrinol. Metab.* 2005;90:700–706. [[PubMed](#)]

60. Smith PN, Theodorakis CW, Anderson TA, Kendall RJ. Preliminary assessment of perchlorate in ecological receptors at the Longhorn Army Ammunition Plant (LHAAP), Karnack, Texas. *Ecotoxicology.* 2001;10:305–313. [[PubMed](#)]

61. Landrum M, Canas JE, Coimbatore G, Cobb GP, Jackson WA, Zhang B, Anderson TA. Effects of perchlorate on earthworm (*Eisenia fetida*) survival and reproductive success. *Sci. Total. Envir.* 2006;363:237–244. [[PubMed](#)]

62. Park J, Rinchar J, Liu F, Anderson TA, Kendall RJ, Theodorakis CW. The thyroid endocrine disruptor perchlorate affects reproduction, growth, and survival of mosquitofish. *Ecotoxicol. Environ. Safety.* 2006;63:343–352. [[PubMed](#)]

63. Patino R, Waincott MR, Cruz-Li EI, Balakrishnan S, McMurry C, Blazer VS, Anderson TA. Effects of ammonium perchlorate on the reproductive performance and thyroid follicle histology of zebrafish. *Environ. Toxicol. Chem.* 2003;22:1115–1121. [[PubMed](#)]

64. Goleman WL, Urquidi LJ, Anderson TA, Smith EE, Kendall RJ, Carr JA. Environmentally relevant concentrations of ammonium perchlorate inhibit development and metamorphosis in *Xenopus laevis*. *Environ. Toxicol. Chem.* 2002;21:424–430. [[PubMed](#)]

65. McNabb FMA. Biomarkers for the assessment of avian thyroid disruption by chemical contaminants. *Avian. Poult. Biol. Rev.* 2005;16:3–10.

66. McNabb FMA, Larsen CT, Pooler PS. Ammonium perchlorate effects on thyroid function and growth in bobwhite quail chicks. *Environ. Toxicol. Chem.* 2004;23:997–1003. [[PubMed](#)]

67. Cheng Q, Perlmutter L, Smith PN, McMurry ST, Jackson WA, Anderson TA. A study on perchlorate exposure and absorption in beef cattle. *J. Agr. Food Chem.* 2004;52:3456–3461. [[PubMed](#)]

68. Aslander A. Experiments on the eradication of Canada thistle, *Cirsium arvarse*, with chlorates and other herbicides. *J. Agric. Res.* 1928;36:915–928.
69. Coates JD, Bruce RA, Patrice J, Achenbach LA. Hydrocarbon bioremediative potential of (per) chlorate-reducing bacteria. *Bioremed. J.* 1999;3:323–334.
70. Van Ginkel GC, Plugge CM, Stroo CA. Reduction of chlorate with various energy substrates and inocula under anaerobic conditions. *Chemosphere.* 1995;31:4057–4066.
71. Wu J, Unz R, Zhang H, Logan B. Persistence of perchlorate and the relative numbers of perchlorate- and chlorate-respiring microorganisms in natural waters, soils and wastewater. *Bioremed. J.* 2001;5:119–130.
72. Batista JR, McGarbey FX, Viera AR. The removal of perchlorate from waters using ion-exchange resins. In: Urbanski T, editor. *Perchlorate in the Environment*. Kluwer Academic/Plenum Press; New York, NY, USA: 2000. Chapter 13.
73. Mihelcic JR. *Fundamentals of Environmental Engineering*. John Wiley & Sons, Inc.; New York, NY, USA: 1999. pp. 119–120.
74. Chiang K, Megonnell M. Ion exchange technologies for perchlorate removal are evolving. *Water World.* 2005;21
75. Cal EPA (California Environmental Protection Agency) Perchlorate Contamination Treatment Alternatives. Draft; Office of Pollution Prevention and Technology Development, Department of Toxic Substances Control; Sacramento, CA, USA: 2004.
76. ITRC (Interstate Technology & Regulatory Council) Perchlorate: Overview of Issues, Status, and Remedial Options. PERCHLORATE-1. Interstate Technology & Regulatory Council, Perchlorate Team; Washington, D.C., USA: 2005. Available online: <http://www.itrcweb.org> (accessed 2006).
77. Polk J, Murray C, Onewokae C, Tolbert DE, Togna AP, Guarini WJ, Frisch S, Del Vecchio M. Case study of ex-situ biological treatment of perchlorate-contaminated groundwater. *Proceedings of 4th Tri-Services Environmental Technology Symposium*; San Diego, CA, USA. 18–20; Jun, 2001.
78. Sartain HS, Craig M. Ex-situ treatment of perchlorate contaminated groundwater. *Proceedings of In Situ and On-Site Bioremediation-The Seventh International Symposium*; Orlando, Florida, USA. 2–5; Jun, 2003.
79. Herman DC, Frankenberger WT. Microbial mediated reduction of perchlorate in groundwater. *J. Environ. Qual.* 1998;27:750–754.
80. Attaway H, Smith M. Reduction of perchlorate by an anaerobic enrichment culture. *J. Ind. Microbiol.* 1993;12:408–412.
81. Wallace W, Ward T, Breen A, Attaway H. Identification of an anaerobic bacterium which reduces perchlorate and chlorate as *Wolinella succinogenes*. *J. Ind. Microbiol.* 1996;16:68–72.
82. Rikken GB, Kroon AGM, van Ginkel GC. Transformation of (per) chlorate into chloride by a newly isolated bacterium: reduction and dismutation. *Appl. Microbiol. Biotechnol.* 1996;45:420–426.
83. Kim K, Logan B. Microbial reduction of perchlorate in pure and mixed culture packed bed bioreactors. *Water Res.* 2001;35:3071–3076. [[PubMed](#)]
84. Bruce RA, Achenbach LA, Coates JD. Reduction of (per) chlorate by a novel organism isolated from paper mill waste. *Environ. Microbiol.* 1999;1:319–328. [[PubMed](#)]
85. Logan BE. Analysis of overall perchlorate removal rates in packed bed reactors. *J. Environ. Eng.* 2001;127:469–471.
86. Hall P. Patented Hall Bioreactor. 2000. Available online: <http://www.perchlorateinfo.com/perchlorate-case-43.html> (accessed August 20, 2007).
87. Van Ginkel GC, Kroon AGM, Rikken GB, Kregen SWM. Microbial conversion of perchlorate, chlorate and chlorite. *Proceedings of the Southwest Focused Groundwater Conference*; National Groundwater Association;

Columbus, OH, USA. 1998. pp. 92–95.

88. Miller J, Logan B. Sustained perchlorate degradation in an autotrophic, gas phase, packed bed bioreactor. *Environ. Sci. Technol.* 2000;34:3018–3022.
89. Logan B, Lapoint D. Treatment of perchlorate and nitrate contaminated groundwater in an autotrophic, gas phase, packed bed bioreactor. *Water Res.* 2002;36:3647–3653. [[PubMed](#)]
90. Evans P, Allyson C, Price S, Moody S, Headrick M, Min B, Logan B. Pilot testing of a bioreactor for perchlorate-contaminated groundwater treatment. *Proceedings of the Third International Conference on Remediation of Chlorinated and Recalcitrant Compounds*; Monterey, California, USA. 20–23; May, 2002.
91. Nerenberg R, Rittmann B, Gillogy T, Lehman G, Adham S. Perchlorate reduction using a hollow fiber membrane biofilm reactor: kinetics, microbial ecology, and pilot-scale studies. *Proceedings of In-situ and On-site Bioremediation-The Seventh International Symposium*; Orlando, FL, USA. 2–5; Jun, 2003.
92. Ping H, Mi Z, Xin-hua X. Research on anaerobic treatment of perchlorate-contaminated water. *Proceedings of the 231st ACS National Meeting*; ACS; Atlanta, GA, USA. 26–30; Mar, 2006.
93. Faris B, Vlassopoulos D. A systematic approach to in situ bioremediation in groundwater. *Remediation.* 2003;13:27.
94. Perlmutter MW, Britto R, Cowan JD, Jacobs AK. In Situ Biotreatment of Perchlorate and Chromium in Groundwater Bioremediation of Inorganic Compounds *Proceedings of 6th International In Situ and On-Site Bioremediation Symposium* Leeson A, Alleman BC, editors. Battelle Press; Columbus, Ohio, USA: 2001. Vol. 6No. 9315–322.322
95. Zawtocki C, Lieberman TM, Birk G. A dash of oil and let marinate. *Pollut. Eng.* 2004;36:30–34.
96. Hatzinger PB. Perchlorate biodegradation for water treatment. *Environ. Sci. Technol.* 2005;39:239A–47A. [[PubMed](#)]
97. EPA (Environmental Protection Agency) Perchlorate treatment technology update. *Federal Facilities Forum Issue Paper.* 2005. Available online: [www.epa.gov/tio/tsp](http://www.epa.gov/tio/tsp) (access 2009).
98. Parette R, Cannon F. The removal of perchlorate from ground water by activated carbon tailored with cationic surfactants. *Water Res.* 2005;39:4020–4028. [[PubMed](#)]
99. Morss C. Perchlorate groundwater treatment. *Pollut. Eng.* 2003;35:18.
100. Liang S, Scott K, Scherzinger R. Perchlorate removal by reverse osmosis, nanofiltration, and ultrafiltration. *Proceedings of the American Water Works Association Membrane Conference*; San Antonio, TX, USA. 4–7; Mar, 2001.
101. Lieberman MT, Zawtocki C, Borden RC, Birk GM. Treatment of perchlorate and trichloroethane in groundwater using edible oil substrate (EOS). *Proceedings of the National Ground Water Association Conference on MTBE and Perchlorate Assessment, Remediation and Public Policy*; Costa Mesa, CA, USA. 3–4; Jun, 2004.
102. Rusanova M, Polaskova P, Muzikar M, Fawcett R. Electrochemical reduction of perchlorate ions on platinum-activated nickel. *Electrochim. Acta.* 2006;51:3097–3101.
103. Lee C, Kramer T. Electrochemical reduction of perchlorate ions on pitting corrosion developed titanium surface. *Proceedings of the 233rd ACS National Meeting*; ACS; Chicago, IL, USA. 25–29; Mar, 2007.
104. Xiong Z, Zhao D. Destruction of perchlorate in fresh and saline water by stabilized zero-valent iron nanoparticles. *Proceedings of the 231st ACS National Meeting*; ACS; Atlanta, GA, USA. 26–30; Mar, 2006.
105. Oh S, Chiu P, Kim B, Cha D. Enhanced reduction of perchlorate by elemental iron at elevated temperatures. *J. Hazard. Mater.* 2006;B129:304–307. [[PubMed](#)]
106. Gurol MD, Kim K. Investigation of perchlorate removal in drinking water sources by chemical methods. In: Urbanski ET, editor. *Perchlorate in the Environment*. Kluwer/Plenum Academic Press; New York, NY, USA: 2000. Chap. 10.

107. Hurley K, Shapley J. Catalytic reduction by hydrogen of perchlorate in water; Proceedings of the 232nd ACS National Meeting; ACS; 10–14; San Francisco, CA, USA: Sep, 2006.

---

Articles from International Journal of Environmental Research and Public Health are provided here courtesy of  
**Multidisciplinary Digital Publishing Institute (MDPI)**